

Automatic Variable Selection

- Instead of choosing what goes into the model manually, we can find a program/algorithm to do so
- Suppose we have p and q predictors. There are 2^q possible regression models
- A computer can potentially pick a "best" model, but it has to be taught what is meant by "best"
- keep in mind: bias-variance tradeoff for prediction based on learned model
 - Bias: the prediction error resulting from miss-specifying the model
 - Variance: the prediction error resulting from variations in the data used for fitting
- Suppose we have a model f to describe response y from feature z . Given a new z_{n+1} we may write

$$y_{n+1} = \underbrace{f(z_{n+1})}_{z_{n+1}^T \beta} + \epsilon_{n+1}$$

- Suppose that we have past data.

$$E[y_{n+1}] = E[z_{n+1}^T \beta] = z_{n+1}^T \beta = E[y_{n+1}] = f(z_{n+1})$$


Namely, the bias is zero.

It may not be zero anymore provided:

We don't have/use an unbiased estimator for f . Typically, this can happen because we don't know precisely f

All Combinations

Example: 3 predictors




$$SS_{res} = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

Summary: .. .

# pred	Best	SS _{res}
0	\emptyset	10
1	{3}	5
2	{1, 2}	2
3	{1, 2, 3}	1

- Forward approach: we start at \emptyset and add the best predictor if the new model is statistically significant; we stop otherwise (using F-test for extra sum of squares)
- Backward approach: we start with all predictors, and drop the least significant / the one that leaves you with **minimal** SS_{res} if the old model is not statistically significantly better than the new one; otherwise, stop
- Issues:
 - = these are greedy approaches (a hybrid approach that "looks" into the future also makes sense)

- Both approaches usually disagree on the stopping point
- We cannot use p-values, confidence intervals, z-stats in the post selection model: "the quiet scandal"
- The distribution of $\hat{\beta}_j$ when we take the model selection process into account:



$$\Pr(\hat{\beta}_j = 0) > 0$$

"The quiet scandal" [Leo Breiman 92]

- why use stepwise approaches:
 - Models can be cross-validated to measure accuracy
 - Always leave out variables which

Can save time & money

Cross Validation

- split the data into k smaller sets; leave one aside; fit a family of models based on the other $k-1$ sets; evaluate accuracy over left-out set; repeat for all $i=1, \dots, k$; average accuracy; pick the model with best averaged accuracy.
- special case $k=n$ called leave-one-out
- Let $\hat{y}_i^{(i)}$ be LS predictor when z_i is left out:

$$\hat{y}_i^{(i)} = z_i^T \hat{\beta}^{(i)}$$

where $\hat{\beta}^{(i)}$ depends on $\{(z_j, y_j)_{j \neq i}\}$
(fitted based on Z with i -th row removed)

- Cross Validation (CV) error:

$$CV(\text{model}) = \sum_{i=1}^n (y_i - \hat{y}_i^{(i)})^2$$

- Why use CV:

- Generally, knocks out models that overfit

- parallels our goal of predicting new values

- Issues: can be computationally prohibitive

CV in Linear Models

- we have a "shortcut"

$$\hat{y}_i = H_{ii}y_i + (1 - H_{ii})\hat{y}_i^{(i)}$$

(proof is based on the Sherman-Morrison formula for the inversion of matrix + rank one mat.)

$$H = Z(Z^T Z)^{-1}Z^T$$

$$\text{so } \hat{y}_i^{(i)} = \frac{\hat{y}_i - H_{ii}y_i}{1 - H_{ii}}$$

all left out prediction are evaluated using one regression instead of n

- The residuals:

$$\frac{y_i - \hat{y}_i^{(i)}}{1 - H_{ii}} = \frac{y_i - \hat{y}_i}{1 - H_{ii}} = \frac{\hat{\epsilon}_i}{1 - H_{ii}}$$

$H_{ii} \leq 1$

- Overall:

$$CV = \sum_i^n (y_i - \hat{y}_i^{(i)})^2 = \sum_{i=1}^n \frac{\hat{\epsilon}_i^2}{(1 - H_{ii})^2}$$

Penalty on number of variables

- Akaike's Information Criterion (AIC)

$$AIC := n \cdot \log\left(\frac{SS_{res}}{n}\right) + 2p$$

- Bayes' Information Criterion (BIC)

$$BIC = n \log \frac{SS_{res}}{n} + p \log n$$

- We pick the model that minimizes AIC or BIC

- Considerations:

-> AIC & BIC are more "principled" approaches than forward/backward

- For $\log(n) > 2$, BIC penalizes predictors more severely.

- BIC is better at getting the "right" model - correctly

identifies non-zero variables when true model is linear as $n \rightarrow \infty$

- AIC is more accurate in making predictions
- We should not use confidence intervals and t-tests after selecting models with AIC or BIC